

THE PROBLEM OF HIPPARCOS DISTANCES TO OPEN CLUSTERS: I. CONSTRAINTS FROM MULTICOLOR MAIN SEQUENCE FITTING¹

Marc H. Pinsonneault

Astronomy Department, Ohio State University,
174 West 18th Avenue, Columbus OH 43210
email: pinsono@astronomy.ohio-state.edu

John Stauffer

Harvard-Smithsonian Center for Astrophysics,
60 Garden St., Cambridge, MA 02138
email: stauffer@cfa.harvard.edu

David R. Soderblom and Jeremy R. King

Space Telescope Science Institute
3700 San Martin Drive, Baltimore MD 21218
email: soderblom@stsci.edu, jking@stsci.edu

Robert B. Hanson

Univ. of California Observatories/Lick Observatory,
University of California, Santa Cruz, CA 95064
email: hanson@ucolick.org

Received _____; accepted _____

¹Based on data from the ESA Hipparcos astrometry satellite.

ABSTRACT

Parallax data from the Hipparcos mission allow the direct distance to open clusters to be compared with the distance inferred from main sequence (MS) fitting. There are surprising differences between the two distance measurements, which indicate either the need for changes in the cluster compositions or reddening, underlying problems with the technique of main sequence fitting, or systematic errors in the Hipparcos parallaxes at the 1 mas level. We examine the different possibilities, focusing on MS fitting in both metallicity-sensitive $B-V$ and metallicity-insensitive $V-I$ for five well-studied systems (the Hyades, Pleiades, α Per, Praesepe, and Coma Ber).

The Hipparcos distances to the Hyades and α Per are within 1σ of the MS fitting distance in $B-V$ and $V-I$, while the Hipparcos distances to Coma Ber and the Pleiades are in disagreement with the MS fitting distance at more than the 3σ level. There are two Hipparcos measurements of the distance to Praesepe; one is in good agreement with the MS fitting distance and the other disagrees at the 2σ level. The distance estimates from the different colors are in conflict with one another for Coma but in agreement for the Pleiades. Changes in the relative cluster metal abundances, age related effects, helium, and reddening are shown to be unlikely to explain the puzzling behavior of the Pleiades. We present evidence for spatially dependent systematic errors at the 1 mas level in the parallaxes of Pleiades stars. The implications of this result are discussed.

Subject headings: Galaxy: Open Clusters and Associations: General; Galaxy: Open Clusters and Associations: Individual (α Per, Coma Ber, Hyades, Pleiades, Praesepe); Stars: Evolution; Stars: Hertzsprung-Russell Diagram

1. The Problem

Main sequence fitting is a basic tool used in the study of star clusters; the principle behind it is also used to estimate distances to field main sequence (MS) stars. The Hipparcos mission (ESA 1997) has provided parallaxes for a number of open cluster stars, which permits a direct determination of the distances to the open clusters which can be compared with distances obtained from MS fitting. There are surprising differences between distances obtained with these two methods; in this paper we explore possible explanations for them.

MS fitting relies upon the Vogt-Russell theorem: the location of a star in the HR diagram is uniquely specified by its mass, composition, and age. This implies that we can infer the distance of a given cluster by comparing the apparent magnitudes of cluster stars with the absolute magnitudes of stars with known composition and distance. There are several possible approaches. Unevolved lower MS field stars with known distances or a cluster (such as the Hyades) of known distance can be used to construct an empirical MS. The distance to the cluster is inferred from the vertical shift needed to line up the cluster MS with the empirical MS. Clusters can also be compared with theoretical isochrones calibrated on the Sun; the latter method requires a color calibration which relates the model effective temperatures to the observed colors.

Most nearby open clusters are close to the Sun in metal abundance, which minimizes uncertainties in the distance scale from variations in composition. There is also a large database of fundamental effective temperature measurements for stars near the solar $[\text{Fe}/\text{H}]$, so the color calibrations should be relatively reliable. The nearby open clusters also have been extensively studied for membership, photometry, abundances, and reddening. For all of these reasons the open cluster distance scale has not been regarded as controversial, and evidence that MS fitting yields incorrect distances could have significant astrophysical

importance.

The Hipparcos mission has resulted in a large increase in the number of open cluster stars with measured parallaxes. This data allows the distance scale inferred from MS fitting to be compared with the distance scale inferred from trigonometric parallaxes. The recently announced Hipparcos determination of the mean parallax of the Pleiades cluster gives the result 8.61 ± 0.23 milliarcsec (van Leeuwen & Hansen Ruiz 1997a). This corresponds to a distance of 116 ± 3 pc, or a distance modulus of 5.32 ± 0.06 magnitude. Traditional determinations of the Pleiades distance (e.g., VandenBerg & Bridges 1984; Soderblom et al. 1993), comparing the cluster’s main sequence to that of nearby stars, lead to a distance modulus of about 5.6 mag ($d \sim 130$ pc; $\pi \sim 7.7$ mas). Thus the Hipparcos parallax, being almost 1 mas larger than expected, suggests that the Pleiades cluster stars are systematically ~ 0.3 magnitude fainter than we have thought up to now.

Parallaxes for stars in other clusters have also been measured, and the results are compared with those obtained from MS fitting in Table 1 (data taken from Perryman et al. 1997, Mermilliod et al. 1997, Robichon et al. 1997). The standard reddening for the clusters is also indicated, along with a notation about whether or not differential reddening is present. The second column lists the cluster [Fe/H] values from Boesgaard & Friel (1990) and Friel & Boesgaard (1992); we have adopted their abundance scale for the clusters in the present study (see Section 4). Mermilliod et al. 1997 and Robichon et al. 1997 concluded that there is no simple explanation for the discrepancies between the MS fitting and Hipparcos distances, and that all of the possible classes of solutions appeared unsatisfactory.

We note that a second calculation of the distance to Praesepe has been performed by van Leeuwen & Hansen Ruiz (1997b), and they find a distance modulus of 6.49 ± 0.15 - in disagreement both with MS fitting and the Mermilliod et al. Hipparcos distance. For the

Table 1. Open Cluster Parameters

Cluster	[Fe/H]	$m - M$ Apparent	$(m - M)_o$ Lynga	$(m - M)_o$ Hipparcos	$(m - M)_o$ This paper	$E(B - V)$ mag
Hyades	+0.13	3.01	3.01	3.33±0.01	3.34±0.04	0.00
Coma Ber	−0.07	4.49	4.49	4.73±0.04	4.54±0.04	0.00
Pleiades	−0.03	5.61	5.48	5.33±0.06	5.60±0.04	0.04
IC 2602		6.02	5.89	5.84±0.07	...	0.04
IC 2391		5.96	5.92	5.83±0.08	...	0.01
Praesepe	+0.04	5.99	5.99	6.24±0.12	6.16±0.05	0.00
α Per	−0.05	6.36	6.07	6.33±0.09	6.23±0.06	0.10 ^a
Blanco 1		6.97	6.90	7.01±0.26	...	0.02
IC 4756		8.58	7.94	7.30±0.19	...	0.20 ^a
NGC 6475		7.08	6.89	7.32±0.19	...	0.06
NGC 6633		8.01	7.47	7.32±0.34	...	0.17 ^a
Stock 2		8.62	7.41	7.50±0.32	...	0.38 ^a
NGC 2516		8.49	8.07	7.71±0.15	...	0.13
NGC 3532		8.53	8.40	8.10±0.36	...	0.04

^aVariable reddening

purposes of this paper we have adopted the Mermilliod distance; if we were to adopt the VH97b distance to the cluster we would have to add Praesepe to the list of clusters with a significant (2σ) discrepancy between the MS fitting and Hipparcos distance scales.

The first column of distance moduli in Table 1 lists the values cited as “Lynga” by Mermilliod et al. (1997) and Robichon et al. (1997). We note that these are *apparent* distance moduli, needing considerable (up to 1.2 mag) corrections for extinction, and cannot be directly compared with the distance moduli $(m - M)_o$ calculated from the Hipparcos parallaxes. The second column in Table 1 lists the distance moduli which correspond to the cluster distances given in Lynga’s (1987) Catalogue. These distances come from a variety of sources, are still scaled to a Hyades distance modulus of 3.01 mag, and need corrections for each clusters metallicity. One motivation for our study is to place MS fitting distances for open clusters on a consistent scale. In a paper in preparation, we have found that the MS fitting distances to some of the more distant open clusters are substantially different from the Lynga distances and in marked disagreement with the Hipparcos parallax distances. A second question is the precision of MS fitting estimates; we will show that accuracy at the 0.05 mag level is possible for well-studied systems. Our results for the clusters studied in this paper are in the fourth column.

Discrepancies between the MS fitting distances and the Hipparcos distances could arise from several sources. As indicated above, one possibility is that the MS fitting distances need to be rederived on a consistent scale. Another possibility is that some of the basic properties of well-studied open clusters, such as composition, age, or reddening, need to be revised. If neither of these possibilities can reconcile the distance scales, then we are left with one of two important conclusions : either there are fundamental problems with MS fitting or there are unrecognized systematic errors in the Hipparcos parallaxes themselves.

These issues are important for other questions as well. For example, recent proposed

revisions to the globular cluster distance and age scales, based on Hipparcos parallaxes of subdwarfs, rely on the same MS fitting technique that gives rise to the puzzling distances to open clusters (Reid 1997; Gratton et al. 1997; Chaboyer et al. 1998; but see also Pont et al. 1998).

In this paper we address the essential issues raised above. The Pleiades, Praesepe, and α Per are well-suited for a more detailed examination. There is good membership information and multicolor photometry for all three; α Per is a system with an age comparable to that of the Pleiades (50 Myr vs. 100 Myr) and therefore it provides a test of age-related effects. We have also examined the Coma Ber star cluster, which has a low quoted error for its Hipparcos distance. In a companion paper (Soderblom et al. 1998) we have searched for field stars with accurate parallaxes and anomalous positions in the HR diagram.

We begin by describing the theoretical models which we use and the open cluster data in section 2. In section 3 we begin with a comparison of the Pleiades, Praesepe, and α Per in different colors. We then use the Hyades cluster to test the zero-point of our distance scale, check on the shape of the isochrones in the observational color-magnitude diagram, and to determine the sensitivity of distance estimates in different colors to changes in metal abundance. We then derive distance modulus estimates at both solar $[\text{Fe}/\text{H}]$ and the individual abundances inferred from high-resolution spectroscopy for the Pleiades, Coma Ber, Praesepe, and α Per using several different methods and both $B-V$ and $V-I$. The Pleiades and Coma Ber are found to be in disagreement with the Hipparcos distance scale. We discuss the sensitivity of our results to age, composition, and reddening in section 4, and present evidence that the Hipparcos parallaxes may contain small-scale (~ 1 deg) systematic effects ~ 1 mas in size, large enough to cause the Pleiades parallax discrepancy. Our conclusions are in section 5.

2. Method and Data

2.1. Theoretical Model Parameters

Theoretical stellar models were constructed with the Yale stellar evolution code for a range of compositions. The nuclear reaction cross-sections are from Bahcall, Pinsonneault, & Wasserburg (1995). We use the Saumon, Chabrier, & van Horn (1995) equation of state for temperatures less than 10^6 K; this EOS is superior to the treatment in earlier versions of the Yale evolution codes for the conditions present in low mass stars. The fully ionized EOS in the Yale code was used for higher temperatures; in numerical tests this produced a MS indistinguishable from that obtained with the OPAL EOS (Rogers, Swenson, & Iglesias 1996).

Model atmospheres from Kurucz (1991a) were used as a surface boundary condition; these are available for a range of metal abundances. We also constructed models (for solar composition only) using a grid of atmospheres kindly provided by F. Allard (see Allard et al. 1997 for a review). Kurucz (1991b) molecular opacities were used for temperatures less than 10^4 K, and OPAL opacities (Rogers & Iglesias 1992) were used for higher temperatures. The mixing length ($\alpha = 1.735$) and helium abundance ($Y = 0.2704$) were calibrated by requiring that a solar model have the radius and luminosity of the Sun at the age of the Sun.

We then generated models from $[\text{Fe}/\text{H}] = -0.3$ to $[\text{Fe}/\text{H}] = +0.2$ in 0.1 dex increments. At each metal abundance we ran a grid of masses from 0.2 - $1.6 M_{\odot}$ in $0.1 M_{\odot}$ increments. Helium abundances for non-solar $[\text{Fe}/\text{H}]$ were obtained by combining the solar helium abundance with an assumed $\Delta Y / \Delta Z = 2$. Isochrones were constructed at a variety of ages, to account for the pre-MS nature of the lower mass stars in the young clusters. We also constructed a set of models with solar Z but enhanced helium ($Y=0.37$). The Yale color

calibration (Green 1988) was used to convert from the theoretical (L , T_{eff}) plane to the observational (magnitude, color) plane in $B-V$ and $V-I$.

2.2. Field Star and Cluster Data

For the open clusters the distance modulus estimates from Hipparcos parallaxes are taken from the Mermilliod et al. (1997) and Robichon et al. (1997) papers. To perform MS fitting we restricted the sample to stars with $B-V$ in the range 0.5 - 0.9. Distance modulus estimates for early-type stars can be affected by the assumed cluster age, and $V-I$ becomes a poor temperature indicator; we therefore excluded stars with $B-V$ colors less than 0.5. The color calibrations become unreliable for late-type stars; we therefore excluded stars with $B-V$ colors greater than 0.9 from our main sequence fitting estimates. Our choice of this color interval also makes our distance estimates insensitive to the adopted cluster ages.

For the photometry of open cluster stars, data was taken from several sources. For the Hyades we used cluster members (as determined by Perryman et al. 1997) which were considered to be single stars by Griffin et al. (1988). We restricted the sample to stars with RI photometry (on the Johnson system) from Mendoza (1967). Since the Hipparcos sample is magnitude-limited, including other stars in common between the Perryman and Griffin paper did not result in increasing the sample size significantly in the color range of interest, and including additional stars with RI colors measured in other systems raises the possibility of systematic color effects. Individual parallaxes for the Hyades stars were used to infer individual M_V values and correct for depth effects.

For both the Pleiades and Praesepe, photometry for the upper MS stars was taken from Mendoza (1967). Additional data for α Per is from Stauffer et al. (1985), Stauffer et al. (1989), Prosser (1992), Mitchell (1960), and Prosser (1994b). Photometry for the

lower MS stars in the Pleiades was taken from Stauffer (1980,1982a,1984), Johnson & Mitchell (1958), and Landolt (1979). Additional photometry for Praesepe was also taken from Upgren, Weis, & DeLuca, (1979), Weis (1981), Stauffer (1982b), and Johnson (1952). Photometry for the Coma cluster was taken from Johnson & Knuckles (1955) and Mendoza (1967).

The Mendoza open cluster RI photometry is on the Johnson system, while the other open cluster RI data is on the Kron system; both the isochrones and the field star data are on the Cousins system.

For the reddening we adopted $E(B - V)=0$ for the Hyades (Crawford & Perry 1966) and Praesepe (Crawford & Barnes 1969). For the Pleiades we adopted $E(B - V)=0.04$ and used individual reddenings for a small number of highly reddened stars (Crawford & Perry 1976, Breger 1986, Stauffer & Hartmann 1987). α Per has patchy and variable differential reddening which is apparent in the cluster color-magnitude diagram; we adopted $E(B - V)=0.10$ (see Crawford & Barnes 1974, Prosser 1994a). We corrected the V magnitudes and different colors for reddening as follows (Allen 1973, Bessell & Brett 1988):

$$A_V = 3.12E(B - V)$$

$$E(V - I)_{Cousins} = 1.25E(B - V)$$

$$E(V - I)_{Kron} = 1.5E(B - V)$$

$$E(V - I)_{Johnson} = 1.75E(B - V)$$

The impact of reddening on distance modulus estimates in different colors is discussed in Section 4. The reddening-corrected $(V - I)_J$ and $(V - I)_K$ were converted to Cousins $(V - I)_C$ using the transformations in Bessell (1979) and Bessell & Weis (1987) respectively:

$$(V - I)_C = 0.778(V - I)_J, 0 < (V - I)_J < 2$$

(true for all Mendoza stars in this color range); and

$$(V - I)_C = 0.227 + 0.9567(V - I)_K + 0.0128(V - I)_K^2 - 0.0053(V - I)_K^3$$

3. Main Sequence Fitting

Most work on cluster distances has used $B - V$ colors as an effective temperature index. In Figure 1 we compare the Pleiades to α Per and Praesepe at the Hipparcos distances in the observational (M_V versus $B - V$) plane. Both Praesepe and α Per are distinctly above the Pleiades. This result is as disturbing as the discrepancies in Table 1 because the measured $[\text{Fe}/\text{H}]$ of the three clusters are within 0.1 dex, implying that the cluster main sequences should be very close in this diagram (within 0.1 magnitudes).

The $B - V$ color is highly metallicity sensitive, and the distances inferred from $B - V$ are therefore quite sensitive to the adopted cluster $[\text{Fe}/\text{H}]$ values. If the true cluster abundances deviate from the currently accepted values then one might expect the cluster main sequences to be closer in a less metallicity-sensitive index, such as $V - I$. The clusters are compared in $(V - I)_{\text{Cousins}}$ in Figure 2. Praesepe is closer to the Pleiades in this index, which suggests that part of the difference in Figure 1 is caused by a difference in chemical composition. However, the two cluster main sequences are still well separated in $V - I$ and the difference between the Pleiades and α Per is the same in both colors. These figures illustrate both the magnitude of the problem and the difficulty in explaining it by either metallicity or age. To quantify this problem, we need to determine the sensitivity of distances based upon temperature measurements from $B - V$ and $V - I$ to changes in composition.

3.1. Theoretical Isochrones and Field MS Data

We show theoretical isochrones for 1 Gyr populations with different abundances in Figure 3. The top, middle, and bottom panels are the theoretical plane, $V - I$, and $B - V$ respectively; the Yale color calibration was used for the bottom two panels. The width of the MS is different in each; 0.5 dex in $[\text{Fe}/\text{H}]$ produces a range of ~ 0.3 mag in the theoretical plane, ~ 0.45 dex in $V/V - I$, and ~ 0.6 dex in $V/B - V$. Helium variations affect the isochrones the same in all three planes : a 0.1 increase in Y produces a 0.28 magnitude decrease in the locus of the main sequence. The isochrones are nearly parallel across the entire color range of interest. The Alonso, Arribas, & Martinez-Roger (1996) [AAR] color calibration is an alternate method of converting from the theoretical to the observational plane. The AAR color calibration is based on application of the Infrared Flux Method, and it can be used to derive the sensitivity of different color indices to changes in metal abundance. The $[\text{Fe}/\text{H}]$ sensitivity of $B - V$ in AAR is comparable to that in the Yale color calibration, but AAR find that $V - I$, at least in the Johnson system, is metallicity independent : that is, that the changes in a $V - I$ based color magnitude diagram should faithfully reflect changes in the theoretical HR diagram. The Hyades and Praesepe clusters provide support for the AAR findings on the metallicity sensitivity of $V - I$, at least for systems near solar metal abundance.

If we adopt the Yale color calibration, a 0.1 dex increase in $[\text{Fe}/\text{H}]$ produces a 0.12 magnitude decrease in M_V at fixed $B - V$ and a 0.09 magnitude decrease in M_V at fixed $V - I$. For example, this would imply that the Hyades (at $[\text{Fe}/\text{H}] = +0.15$) would lie 0.18 magnitudes above a solar composition isochrone in $B - V$ and 0.135 magnitudes above a solar composition isochrone in $V - I$. If the metallicity sensitivity of AAR is adopted, then a 0.1 dex increase in $[\text{Fe}/\text{H}]$ produces a 0.13 magnitude decrease in M_V at fixed $B - V$ and a 0.06 magnitude decrease in M_V at fixed $V - I$. The Yale color calibration implies that

a color-color diagram in $B-V$ and $V-I$ should be metallicity-insensitive because both indices are metallicity sensitive (albeit to slightly different degrees); the AAR results would produce a wider spread in a color-color diagram because $B-V$ is much more metallicity sensitive than $V-I$.

The local field stars span a range in $[\text{Fe}/\text{H}]$, with the bulk of the F and G stars in the Gliese catalog spanning a range from -0.3 to +0.2 (Wyse & Gilmore 1995; see also McWilliams 1997). We therefore expect a significant intrinsic width to the field star MS as well as a population of binary stars above the MS. The relative width of the MS in the $B-V$ and $V-I$ planes can be used as a test of the relative sensitivities of the two color indices to $[\text{Fe}/\text{H}]$.

The Hipparcos mission has provided parallaxes accurate to 5% or better for a significant number of lower MS stars. There is a smaller sample (680 stars) with both $B-V$ and $V-I$ colors; the field MS in both colors is compared with the models in Figure 4. The field MS is much tighter in $V-I$ than in $B-V$, suggesting that much of the spread in Figure 4 is caused by atmospheric effects (the dependence of $B-V$ color on metal abundance) rather than by interiors effects. We therefore expect that there will be systematic differences between the derived distances in $B-V$ and $V-I$ if the adopted $[\text{Fe}/\text{H}]$ for the isochrone departs from that of the star, or cluster, at the 0.07 (AAR) or 0.03 (Yale) magnitude level per 0.1 dex in $[\text{Fe}/\text{H}]$. This discrepancy will be in the sense that the $V-I$ distance will be longer than $B-V$ if the true metal abundance is higher and the $V-I$ distance will be shorter than the $B-V$ distance if the true metal abundance is lower.

3.2. The Hyades

The Hyades cluster provides an opportunity to check the distances derived from MS fitting against the Hipparcos distance scale for a system with a large number of measured parallaxes spread across a large region of the sky. We can also compare the isochrones in the theoretical plane and in different colors; the Hyades provides a useful check of the sensitivity of the isochrones to changes in metal abundance because it has a metal abundance 0.1-0.2 dex above solar.

The Perryman et al. (1997) paper provides locations for Hyades stars in the theoretical plane as well as isochrones for both solar composition and the Hyades $[\text{Fe}/\text{H}]$ adopted in that study, 0.14 ± 0.05 . Our 600 Myr isochrones for both $[\text{Fe}/\text{H}]=0$ and $[\text{Fe}/\text{H}]=+0.14$ are compared with the Perryman et al. isochrones in Figure 5. For the range of $3.68 - 3.84$ in $\log T_{\text{eff}}$ our $[\text{Fe}/\text{H}]=0$ isochrone is on average 0.044 magnitudes brighter than the Perryman et al. isochrones. By comparison, a zero-age MS for $[\text{Fe}/\text{H}]=0$ provided by Vandenberg (private communication) is 0.032 magnitudes fainter than our 100 Myr isochrone and for $\log T_{\text{eff}} > 3.75$ the Yale and Vandenberg isochrones agree to within 0.008 magnitudes. This comparison indicates that systematic differences between different isochrones are at or below the 0.04 magnitude level overall and agree to within 0.03 magnitudes near the temperature of the Sun.

The Hyades MS of Perryman et al. is 0.164 magnitudes brighter than their solar $[\text{Fe}/\text{H}]$ isochrone in this temperature interval; their isochrone with $[\text{Fe}/\text{H}]=+0.14$ and solar scaled helium would be too faint to be consistent with the data. They were therefore forced to a subsolar helium abundance (0.26) to reproduce the observed Hyades MS. Our $[\text{Fe}/\text{H}]=+0.14$ isochrone with $Y = 0.283$ is 0.017 magnitudes fainter than the Hyades MS from Perryman et al.; this implies that our models are consistent with the Hyades having the $[\text{Fe}/\text{H}]$ inferred from high-resolution spectroscopy and $\Delta Y/\Delta Z=2$. This result is obtained largely because

our solar composition isochrone is slightly brighter than the Perryman et al. isochrone; we constructed isochrones with different helium and verified that the changes in the position in the theoretical HR diagram resulting from changes in Y and Z agree with the offsets in the Perryman et al. paper to better than 5%. We obtain similar results when fitting in the observational HR diagram using $B-V$ and $V-I$. This illustrates the importance of small effects when inferring helium abundances based upon HR diagram position.

The absolute V magnitudes of single stars and binaries in the Hyades are shown as a function of $B-V$ and $V-I$ in Figure 6. Half of the stars are binaries and the binaries scatter systematically above the single stars in the color-magnitude diagram. For comparison, the Schwan (1991) empirical MS for the Hyades in $B-V$, shifted to a distance of 3.33, is shown. We also derived an empirical fit to the single star sequence in the $V-I$ plane which is compared with the cluster data. The Hyades abundance isochrones are shifted up in the HR diagram by 0.18 magnitudes in $B-V$ and 0.135 magnitudes in $V-I$ relative to a solar abundance isochrone for the metallicity sensitivity in the Yale color calibration. If the metallicity sensitivity of Alonso et al. (1996) is adopted, the Hyades isochrones are 0.19 magnitudes above the solar $[\text{Fe}/\text{H}]$ isochrones in $B-V$ and 0.09 magnitudes above the solar $[\text{Fe}/\text{H}]$ isochrones in $V-I$. A slight mismatch between the shape of the isochrones and the empirical MS is present, and the distance modulus estimates from the isochrones are clearly close to the Hipparcos distance scale.

For each of the 35 single stars in our sample we derived a distance modulus estimate; the average is 3.36 with a RMS deviation of 0.16. Since the average error in M_V is 0.13, color errors are contributing little to the overall scatter in the diagram. Since the parallax errors can be correlated, the error in the mean scales as $N^{-0.35}$, not $N^{-0.5}$ (Lindgren 1988, 1989); the formal error in the mean distance modulus estimate is therefore 0.05. Because of the nearness of the Hyades the mean distance modulus estimate will depend on the subset

of the stars used in the comparison, and therefore the difference between this estimate and the cluster mean of 3.33 ± 0.01 is not problematic.

The difference ($M_V(\text{observed}) - M_V(\text{predicted})$) for both colors is shown for the single stars as a function of color in Figure 7; the difference between the isochrones and the empirical MS is also shown. For the Yale color calibration the mean for $B - V$ is -0.04, implying a distance modulus of 3.32; the mean for $V - I$ is +0.05, implying a distance modulus of 3.41. The dispersion about the mean in both cases is 0.13, consistent with the errors in the absolute magnitudes.

The discrepancy between the distance estimates in $B - V$ and $V - I$ could be reduced, or removed, by an increase in the adopted cluster metal abundance; alternately, it could indicate that $B - V$ is more metallicity sensitive and $V - I$ is less metallicity sensitive than predicted by the models. Since increasing the Hyades metal abundance would cause a disagreement with both the parallax distance to the cluster and the spectroscopic $[\text{Fe}/\text{H}]$ measurements we believe that the latter explanation is more likely. If we adopt the derivatives of M_V with respect to $[\text{Fe}/\text{H}]$ from AAR the mean for the $B - V$ distance is 3.33 (-0.03) and the mean for the $V - I$ distance is 3.36 (0.0). The distance we obtain by averaging the two colors is 3.34, in excellent agreement with the Hipparcos distance. *We therefore adopt the Yale color calibration at solar metal abundance to set the zero point of the distance scale and the shape of the isochrones and adopt the metallicity sensitivity of the Alonso et al. (1996) color calibration: a 0.1 dex increase in $[\text{Fe}/\text{H}]$ produces a decrease in M_V of 0.13 magnitudes at fixed $B - V$ and 0.06 magnitudes at fixed $V - I$.*

We have added the binaries and binned the deviations between the individual distance modulus estimates and those predicted from the isochrones. The resulting histograms are plotted in Figure 8. Binaries scatter to systematically lower distance estimates relative to single stars; $V - I$ is more affected by this than $B - V$. There are two methods that

can be used to account for binaries. There is a clear separation between the small mass ratio binaries and the single stars and high mass ratio binaries; the peak in the histogram is resistant to the presence of binaries, although there is a slight bias to lower distance modulus estimates in $V - I$ because of its higher sensitivity to binaries. The peaks in the histograms of Figure 8 are at -0.05 and 0 for $B - V$ and $V - I$ respectively, which are in good agreement with the single star averages of -0.03 and 0. Alternately, the distance modulus estimates can be ranked, stars more than 0.2 mag above and below the histogram peak excluded, and the median distance modulus estimate can be inferred. Medians inferred with this technique are also -0.05 and 0.0 for $B - V$ and $V - I$ respectively.

We also compare the distance estimates derived from the isochrones and the distance estimates derived from the empirical Hyades MS with the zero point adjusted to agree with the isochrones at one solar mass for the different colors. The Hyades MS needs to be corrected for the higher than solar $[\text{Fe}/\text{H}]$ of the cluster; the M_V at fixed color for a solar abundance MS is 0.19 magnitudes higher at fixed $B - V$ and 0.09 mag higher at fixed $V - I$. The empirical MS that we adopted at solar metal abundance in $B - V$ and $V - I$ respectively are therefore

$$M_V = -2.75 + (4.03 + 85.7(B - V))^{0.5} \text{ and}$$

$$M_V = -1.976 + 13.758 * (V - I) - 5.427 * (V - I)^2 \text{ (valid only from 0.55 to 0.9 in } V - I\text{).}$$

For younger clusters the empirical Hyades MS needs to be corrected for age effects (a 0.04 magnitude level effect for the Pleiades). We took the difference between our 100 and 600 Myr isochrones and applied it to the relationships given above for the Pleiades. We stress that the distances obtained in this manner have the same zero-point as the isochrones; we are using the shape of the Hyades MS and not its absolute distances, and our distance scale is therefore not tied to the distance to the Hyades (although it is in agreement with

the cluster distance as measured by Hipparcos).

3.3. The Pleiades

In Figure 9 we present histograms of the distance modulus estimates for the Pleiades using different techniques. In the top panels 100 Myr solar composition isochrones were used to estimate M_V from $B-V$ and $V-I$ respectively; in the bottom panels the empirical Hyades MS was used. The darker bins are for hotter stars and the lighter bins are for cooler stars; a discrepancy between the mean of the two is an indication of a deviation between the shape of the isochrone and the cluster CMD. In the isochrone distances cooler stars give systematically longer distance modulus estimates at the 0.1 mag level; these color trends are removed relative to the Hyades MS fits in the lower panels. The good agreement between different techniques suggests that there are small internal errors in MS fitting for systems with good photometry.

An average of the $V-I$ distance methods yields 5.63 ± 0.02 , while an average of the $B-V$ distance methods also gives 5.63 ± 0.02 . This should be compared with the Hyades, where a solar abundance isochrone would give distance modulus estimates that differ by 0.1 magnitudes; this difference is caused by a cluster $[\text{Fe}/\text{H}]$ 0.15 dex higher than solar. If we add the errors in quadrature this implies that we have a 0.03 magnitude relative error in the $B-V$ and $V-I$ distance estimates, which corresponds to a 0.05 magnitude error in the photometric $[\text{Fe}/\text{H}]$. Boesgaard & Friel (1990) obtained $[\text{Fe}/\text{H}] = -0.034 \pm 0.024$ for the Pleiades; at this metal abundance our $B-V$ and $V-I$ distance moduli are 5.59 and 5.61 respectively. Our MS fitting distance to the Pleiades is therefore 5.60, and we estimate that errors in the metal abundance and systematic differences in the MS fitting technique are at the 0.04 magnitude level.

In Figure 10 we compare the Pleiades to a $[\text{Fe}/\text{H}]=-0.03$ 100 Myr isochrone in both $B-V$ and $V-I$. The isochrone has been shifted to a distance of 5.60. The isochrone is an excellent fit to the cluster CMD. We note that single rapid rotators are on or above the MS in $V-I$ but below it in $B-V$; this may indicate that the relationship between color and temperature for these stars is different than for slow rotators, and we therefore exclude them from distance estimates in both this cluster and α Per.

A detailed binary inventory for the Pleiades has recently been published (Bouvier, Rigaut, & Nadeau 1997; see also Mermilliod et al. 1992), and we can therefore check for the possible impact of binary contamination on our distance estimates. We compare distance estimates for binaries, single stars, and rapid rotators in Figure 11. As expected, stars with very low distance estimates are binaries and for the high mass ratio binaries $B-V$ is less sensitive than $V-I$. The broader distributions for $V-I$ seen in Figure 9 are therefore a reflection of its greater sensitivity to binary contamination. The techniques that we have applied do not appear to be affected significantly; if we use only the single stars a slightly higher distance of 5.65 is indicated for both colors.

To reproduce the Hipparcos distance of 5.33 we would require $[\text{Fe}/\text{H}] = -0.25$ for $B-V$ and $[\text{Fe}/\text{H}] = -0.45$ for $V-I$. Reproducing the Hipparcos distance to the Pleiades by changing the metal abundance would therefore require a much lower metallicity than that obtained by high-resolution spectroscopy; furthermore, the distance estimates for different colors would be in strong disagreement. Other possibilities are discussed in Section 4.

3.4. Praesepe

In Figure 12 we present histograms of the distance modulus estimates for Praesepe using different techniques; as for the Pleiades the top panels are relative to the isochrones

(600 Myr for Praesepe) and the bottom panels are relative to the isochrones with the shape adjusted to agree with the empirical Hyades MS. Particularly for $B-V$, there is a clear indication that the distance estimates for the uncorrected isochrones depend on color at the 0.1 magnitude level; this is more apparent for Praesepe than for the Pleiades largely because the Praesepe sample includes more cool stars than the Pleiades sample. However, the histogram peaks and medians are similar for the uncorrected isochrones and those obtained relative to the shape of the Hyades MS. An average of the $V-I$ distance methods yields 6.17 ± 0.02 , while an average of the $B-V$ distance methods yields 6.08 ± 0.02 . This difference is significantly larger than our estimated relative error of 0.03. We therefore conclude that the difference between the two is real and indicates that Praesepe is mildly metal-rich.

The two distance modulus estimates agree at $[\text{Fe}/\text{H}]=+0.13$ and a distance modulus of 6.25. The Friel & Boesgaard (1992) $[\text{Fe}/\text{H}]$ for Praesepe is $+0.04 \pm 0.04$, while a higher abundance is inferred by some other studies (see Section 4.2). At the FB92 $[\text{Fe}/\text{H}]$ the distance modulus is 6.13 in $B-V$ and 6.19 in $V-I$; we therefore adopt a distance modulus estimate of 6.16 for Praesepe (Figure 13). This is well within the error bounds of the Hipparcos distance estimate of 6.24 ± 0.12 . We conclude that Praesepe is consistent within the errors with the Hipparcos distance measurement and that the photometry is consistent with it being more metal-rich (at the 0.1 dex level) than the FB92 estimate.

The dominant source of error in the distance modulus is the metal abundance of the clusters; in general, relative metal abundances can be determined more precisely than absolute metal abundances. Another way of looking at the problem of reconciling the MS fitting and Hipparcos distance scales is therefore to look at relative distances in different colors and asking what metal abundance difference is needed to explain the results. The Pleiades and Praesepe are especially difficult to explain in combination. At solar $[\text{Fe}/\text{H}]$

the relative distances of these two clusters from MS fitting are 0.45 and 0.54 magnitudes in $B-V$ and $V-I$ respectively. By comparison, the magnitude difference between the two clusters for the Hipparcos distance scale is 0.91. If Praesepe is metal-rich or the Pleiades is metal poor then the true difference in distance modulus estimates will be larger than at solar $[\text{Fe}/\text{H}]$, with $B-V$ being more metallicity sensitive than $V-I$. Reconciling the relative cluster distances in $B-V$ and $V-I$ by changing the metal abundances would require a metallicity difference of 0.35 dex and 0.6 dex respectively; both are well outside the range of relative metal abundances reported by different investigators.

3.5. α Per

α Per is a young system (50 Myr), and it has a larger overall reddening (0.10) than the other clusters we examine and some differential reddening. In Figure 14 we show the distribution in distance modulus estimates in both $B-V$ and $V-I$ relative to a solar composition 50 Myr isochrone. There is a larger population of rapid rotators in this cluster than in the Pleiades, and they show the same pattern (long distances in $B-V$ and short distances in $V-I$). A subpopulation of stars at higher distances is present in both colors, with distances systematically higher for $B-V$ than for $V-I$. This could be caused by variable reddening, rapid rotators with low $\sin i$, or contamination of the sample by non-members. Excluding these stars only affects the distance estimates at the 0.02 magnitude level and does not change the relative distances in the two colors.

There is a well-defined peak in $B-V$ at a distance of 6.275 and the distribution for $V-I$ is centered at the same distance; the median of the single stars is at 6.29 and 6.27 for $B-V$ and $V-I$ respectively, giving average distances of 6.28 and 6.27 for the two colors. Therefore, at solar abundance the average cluster distance is 6.28; there is a hint of a mild metal deficiency in the relative distances in the two colors, at the 0.02 dex level. If we

adopt the high-resolution abundance $[\text{Fe}/\text{H}]=-0.05$ our average distance is 6.23; because the error in the metal abundance is higher for this system than for the others (0.05) the error is larger, 0.06 magnitudes. A 50 Myr isochrone is compared with the cluster in Figure 15. Significantly, there is no evidence for a discrepancy between the MS fitting and Hipparcos distance scales; because both α Per and the Pleiades are young, this indicates that the problem with the Pleiades is not a consequence of systematic color errors arising from the youth of the system.

3.6. Coma Ber

Coma Ber is a sparse cluster with an age comparable to the Hyades; it is mildly metal-poor ($[\text{Fe}/\text{H}]=-0.071 \pm 0.020$; see Taylor 1994). Surveys have been undertaken to find low-mass members, and few candidates have been found (see Randich, Schmitt, & Prosser 1996 for a discussion). We used the Johnson & Knuckles (1955) photometry for $B-V$ and the Mendoza (1967) photometry for $V-I$; we note that the $V-I$ photometry for the cluster stars listed in RSP differs significantly from that in the Mendoza 1967 paper and is based on an earlier study by Mendoza. The sample size for the color interval used in the other clusters is small (15 stars), so we also included 9 additional stars with $B-V$ from 0.35 to 0.49 and $V > 7.5$. A histogram of the distance estimates from the isochrones is shown in Figure 16. There is a clear peak in the histogram for $B-V$ at a distance modulus of 4.625 at solar $[\text{Fe}/\text{H}]$. The Hyades MS shape yields a peak at a similar distance of 4.675, but with a systematic dependence of the distance estimate on color, i.e. the shape of the Coma Ber and Hyades MS are different for the hotter stars. We therefore adopt the isochrone fit for our distance estimate. Correcting for metal abundance we get a $B-V$ distance modulus for Coma Ber of 4.54; given the low quoted error in the cluster $[\text{Fe}/\text{H}]$ we estimate an error of 0.04 magnitudes. If we were to adopt a larger uncertainty of 0.05

dex in the cluster $[\text{Fe}/\text{H}]$ the error in the distance estimates for all of the clusters we have studied would rise to 0.06 mag. The MS fitting distance of 4.54 is a 3.4σ discrepancy with the Hipparcos distance if we adopt an error in the MS fitting distance of 0.04 and 2.6σ if we adopt an error in the MS fitting distance of 0.06.

The behavior of the cluster in $V - I$, however, is puzzling. There is no well-defined peak for the cool stars, and the hotter stars concentrate at a distance (5.1) well above either the $B - V$ or the Hipparcos distance. This can be traced to the temperature scale for the two colors; the $V - I$ colors for the cluster F stars imply temperatures significantly hotter than the $B - V$ colors. The isochrones are brighter at the hotter temperatures, causing higher distance modulus estimates for these stars (smaller M_V implies larger m-M). We compare the temperatures from the isochrones in the two colors to those obtained by Boesgaard (1987) in a study of lithium in F stars in Figure 17. Boesgaard estimated temperatures from $B - V$, Stromgren photometry, and also measured spectroscopic temperatures; her temperature scale is in excellent agreement with the $B - V$ temperature scale in the isochrones and in significant disagreement (at the 200-300 K level) with the $V - I$ temperature scale for the F stars. Adopting the hotter temperature scale implied by the $V - I$ colors would raise a series of problems : the lithium dip in Coma would be at hotter T_{eff} than for other clusters, it disagrees with spectroscopic temperature estimates and those from Stromgren photometry, and large internal variations in the derived iron abundances for cluster stars would result. We have no explanation for this problem, and reobserving the cluster stars in $V - I$ and IR colors would be useful to understand the problem. We therefore do not use the $V - I$ distance to the system, and our MS fitting distance to this cluster must be taken with this discrepancy in mind. The cluster is compared to isochrones in both of our colors in Figure 18.

4. Discussion

The results of Section 3 indicate that it is the Hipparcos distance to the Pleiades which is in the most serious conflict with MS fitting. In all of the other systems except Coma Ber, MS fitting in different colors yields distance results that are consistent with one another, normal helium, and $[\text{Fe}/\text{H}]$ values from high resolution spectroscopy. Coma Ber may have an equally serious disagreement, but the unusual behavior of the cluster in $V - I$ suggests that other problems may be contributing to the discrepancy for it. We therefore examine in turn the various possible mechanisms that could reconcile the cluster distance scales for the Pleiades; in all cases we believe that they cannot do so. In a companion paper we show that the same conclusions result from an examination of nearby field stars (Soderblom et al. 1998). We then proceed to an analysis of the Hipparcos parallaxes for the Pleiades, and show that there are indications of possible systematic errors that could be the origin of the discrepancy.

The calculations that we have presented are standard stellar models. We have therefore not included physical processes such as gravitational settling, rotational mixing, magnetic fields, internal gravity waves, or mass loss, which are surely present. There are strong reasons for believing that these nonstandard effects will not influence the distance scale, although they could be potentially important for other issues. The single most important reason is the youth of the clusters that we have examined; detailed nonstandard calculations predict little, if any, effect for ages as young as the Pleiades. In addition, any such process would have to affect stars with a wide range in masses to a similar extent and be different among different clusters to explain the pattern that we see.

Gravitational settling is minimal in young systems such as the Pleiades, and the degree to which helium and heavy elements sink depends strongly on the convection zone depth and thus the stellar mass. For example, helium and heavy element diffusion are a

10% fractional effect in the Sun, which is almost 50 times older than the Pleiades. The observed cluster lithium abundances require a mild envelope mixing process, and models with rotational mixing that are consistent with the lithium data predict little or no deep mixing (Pinsonneault 1997). In addition, the observed range in rotation rates in clusters is large, and any extra mixing would produce a spread in MS properties rather than a uniform shift in the distance estimates. Other physical processes could affect the results, but they are still subject to a variety of observational constraints which make a large effect unlikely.

We have compared different standard model calculations, and the zero-point offset is small (0.01-0.03 mag for stars between 5600 and 7000 K, for example). The systematic errors in the standard model distance estimates is therefore also too small to explain the results that we have obtained. We now discuss age, composition, and reddening effects.

4.1. Age and Stellar Activity

It is well-known that many young stars are heavily spotted; this could influence the color-temperature relationship and therefore the distance estimates for young systems such as the Pleiades and α Per. In Figures 1 and 2 we compared these two clusters at the Hipparcos distances in our two colors; the Pleiades is clearly anomalous with respect to α Per if the Hipparcos distance scale is adopted. Since α Per is younger and has a larger population of rapid rotators, if anything α Per should be more anomalous than the Pleiades if our MS fitting age estimates were in error because of activity. We note that similar conclusions can be obtained by comparing young and old field stars (Soderblom et al. 1998). The narrow width of the Pleiades MS also indicates that a wide range in stellar activity does not produce a significant effect on the color-temperature relationship. For all of these reasons we reject the idea that youth is responsible for the difference between the distance estimates.

Another possibility is that activity could be influencing the Pleiades $[\text{Fe}/\text{H}]$, which has been derived from LTE model atmospheres. If such a phenomenon were at play, it might lead to derived abundances being a function of line strength due to the direct effect of activity on the stronger lines formed at smaller depths in the photosphere. We have a number of high resolution spectra of Pleiades members that was originally obtained to study lithium abundances. We have analyzed the Fe I data in the cool Pleiades dwarfs and find no such $[\text{Fe}/\text{H}]$ -line strength correlation. This does not exclude such a real correlation, though, given the influence of damping, which is adjusted to enforce such a lack of correlation. To the extent that our damping assumptions seem quite reasonable compared to numerous other fine spectroscopic analyses, and are consistently applied in both the stellar and solar analyses to yield line-by-line $[\text{Fe}/\text{H}]$ values, the analysis suggests any such trends are not substantial. Regardless, any systematic error in the inferred mean $[\text{Fe}/\text{H}]$ is greatly mitigated by the fact that the damping adjustments enforce consistency with the weaker lines, which are formed at deeper depths, and thus presumably are more immune from the direct effects of chromospheric activity. Activity in very young stars can manifest itself in the form of an effective veiling continuum. Such behavior would presumably weaken the line absorption, thus leading to *underestimated* line strengths and, hence, abundances. Detailed NLTE line formation calculations to determine how the active Pleiades dwarfs' Fe and other metal abundances might be affected by activity, spots, convective flows, *etc.* would be of interest, but are unlikely to produce large errors for the reasons discussed above.

4.2. Heavy Metals

4.2.1. The Cluster $[\text{Fe}/\text{H}]$ Scale

Homogeneous Fe abundances are available for the Pleiades, Praesepe, and α Per from the work of Boesgaard and collaborators. Independent modern fine analyses of these clusters

(and a few others) by other investigators are available for comparison with their work. All the studies considered here derive self-consistent solar Fe abundances with which the stellar values are normalized. Such a careful differential procedure can greatly reduce errors introduced by varying assumptions concerning the solar Fe abundance, model atmospheres, gf values, *etc.*

Boesgaard *et al.* (1988) determine a mean Pleiades iron abundance of $[\text{Fe}/\text{H}] = -0.03$ from analysis of 17 F stars. The mean star-by-star reddening they use is essentially identical to the value we have adopted. Boesgaard (1989) determined a “best” Pleiades abundance by analyzing new data for 8 Pleiads; the result was $[\text{Fe}/\text{H}] = +0.02$. Boesgaard & Friel (1990) used new data for 12 of the same stars in Boesgaard *et al.* to find a mean $[\text{Fe}/\text{H}] = -0.03$. The single datum standard deviation in all these studies is ~ 0.07 dex. The 1σ level error in the mean is 0.02-0.03 dex, so the internal statistical uncertainties appear to be small. Cayrel *et al.* (1988) derive a mean Pleiades $[\text{Fe}/\text{H}]$ of +0.13 from analysis of four Pleiades dwarfs, three of which are significantly cooler (mid G) than the Boesgaard F stars. The standard deviation is 0.10 dex, which is somewhat smaller than their estimated individual errors; the error in the mean is ~ 0.06 dex. The ~ 0.1 dex offset between the Cayrel and Boesgaard values is representative of uncertainties in reddening (which enters via photometric T_{eff} determinations by Boesgaard), the T_{eff} determinations (the Cayrel values are based on $\text{H}\alpha$ profiles), and other details. The Cayrel result is consistent with Eggen’s (1986) inference from narrow band photometry that the Pleiades $[\text{Fe}/\text{H}]$ is near the Hyades value

In order to increase the sample of Pleiades stars with $[\text{Fe}/\text{H}]$ determinations, some of us (King *et al.* 1997) have used high quality Keck spectra of two slowly rotating very cool ($T_{\text{eff}} \sim 4500$ K) Pleiades dwarfs to derive Fe abundances. Our T_{eff} values are spectroscopic determinations from balancing the abundances as a function of excitation potential, and

the normalized abundances are derived by comparison with similarly analyzed solar data on a line-by-line basis. The mean abundance is $[\text{Fe}/\text{H}] = +0.06$, with estimated errors in the mean of perhaps 0.05 dex. While comparison of the different studies indicates there may be systematic errors at the 0.1 dex level, we regard this (dis)agreement to be quite satisfactory given the ~ 2000 K range in T_{eff} , the disparate sources of data, and distinct methods used to derive T_{eff} . While a slightly sub-solar Fe abundance is often assumed for the Pleiades based on the Boesgaard & Friel results, the totality of the high-resolution spectroscopic evidence may be more consistent with a slightly super-solar value; our photometric $[\text{Fe}/\text{H}]$ is consistent with solar $[\text{Fe}/\text{H}]$. Therefore, if anything the data suggest a distance modulus estimate larger than our MS fitting value rather than smaller.

Fe abundances for Praesepe F dwarfs have been derived by Boesgaard & Budge (1988), Boesgaard (1989), and Friel & Boesgaard (1992). The resulting values are $+0.14$, $+0.10$, and $+0.05$, with star-to-star scatter of 0.06-0.07 dex, and mean uncertainties of 0.03-0.04 dex; again, the internal precision is good. The zero-reddening assumed in their T_{eff} determinations is identical to our assumption. Other detailed studies of numerous Praesepe stars comparison are lacking. Analysis of the primary component of the Praesepe SB2 KW367, a mid-G star which is significantly cooler than the Boesgaard F stars, by King & Hiltgen (1996) yielded $[\text{Fe}/\text{H}] = +0.01$ with an uncertainty near 0.05 dex. Again, systematic errors at the 0.1 dex are indicated by this limited comparison. Combined with the above results, we see that $[\text{Fe}/\text{H}]$ for Praesepe is 0.00-0.15 dex larger than for the Pleiades, with a preference for the lower middle of this range. The results inferred from MS fitting are consistent with the upper end of the range.

Boesgaard *et al.* (1988), Boesgaard (1989), and Boesgaard & Friel (1990) derived Fe abundances in α Per F stars. The mean $[\text{Fe}/\text{H}]$ values are -0.02 , $+0.00$, and -0.05 . The α Per Fe abundance seems nearly identical to the Boesgaard Pleiades estimate. The

star-to-star scatter in the larger α Per samples is 0.08-0.09 dex; mean uncertainties are ~ 0.04 dex. The mean of the individual α Per reddening values employed by Boesgaard is ~ 0.03 dex lower than the single value adopted here. This difference might require a 0.05-0.10 dex increase in $[\text{Fe}/\text{H}]$ for consistency with our assumptions. Balachandran *et al.* (1988) determined Fe abundances in a very wide range (F to K type) of α Per stars. The mean abundance of the stars not considered by them to be non-members is $[\text{Fe}/\text{H}] = +0.04$ with a star-to-star scatter of 0.14 dex; the mean internal error is only 0.02 dex. Their assumed reddening is identical to our value. The results of Boesgaard *et al.* and Balachandran *et al.* agree to within 0.1 dex, but when adjustment is made for the slightly different reddening assumptions, the agreement is within a few hundredths of a dex if not exact. Our photometric $[\text{Fe}/\text{H}]$ is slightly sub-solar, at the 0.01-0.02 dex level. It thus appears that the Fe abundance of α Per is not significantly larger than for the Pleiades.

In sum, internal errors in the Fe abundances of main sequence Pleiades, Praesepe, and α Per stars derived from careful homogeneous analyses employing high quality data lead to uncertainties of only 0.05-0.10 dex in relative cluster abundances. We have seen that systematic effects due to errors in reddening, differences in the analysis methodology, *etc.* may approach 0.15 dex. These are small compared to the offset needed to explain the Hipparcos-based M_V values for the Pleiades. Barring fundamental failure or incompleteness in our understanding of spectral line formation and stellar atmospheres, the extant data suggests that the Fe abundances of the Pleiades, Praesepe, and α Per are within ~ 0.10 dex of each other. We might caution, however, that the abundances of other important atmospheric opacity contributors (*e.g.*, *Mg* and *Si*) are, unfortunately, unknown.

4.2.2. *Photometric Constraints and the Binary Distance to the Pleiades*

There are other factors that make a large error in the Pleiades $[\text{Fe}/\text{H}]$ unlikely. Colors that incorporate an infrared band are less sensitive to metallicity than $B-V$. The figures in the previous section indicate clearly that the shift in the cluster distance modulus is the same for different color indices; the Pleiades must be intrinsically subluminous if the revised distance estimate is correct. The deviations from the high-resolution $[\text{Fe}/\text{H}]$ values for the Pleiades are both large and inconsistent from color to color. The spectroscopic binary HD 23642 also provides a distance of 5.61 ± 0.26 consistent with MS fitting, albeit with a large error (Giannuzzi 1995).

4.3. CNO Abundances

Carbon, nitrogen, and oxygen can affect stellar structure in ways other elements do not; are they anomalous in the Pleiades? As part of his thesis, King (1993) examined the oxygen abundances of stars in several clusters over a broad range of age. The $[\text{O}/\text{H}]$ for the Pleiades was found to be higher than for Praesepe (+0.29 and +0.02 respectively, with errors in the mean of 0.08 for both). However, the trustworthiness of abundances (such as these) derived from the high excitation 7774 Å O I lines is a matter of some debate. Besides possible large data and analysis differences between various studies (e.g. King & Hiltgen 1996), there may be significant abundance corrections due to non-LTE effects on line formation in stellar atmospheres (see Garcia Lopez et al. 1995). Unfortunately, systematic errors of 0.3 dex in the cluster O abundances derived from high-excitation lines remains plausible. In any case, the King results would act to make the Pleiades more metal-rich and therefore require a higher distance modulus estimate. Detailed abundance studies would be useful, but deviations from the solar mixture would need to be very large to have a significant impact on the luminosity of the MS.

4.4. Helium

The initial solar helium abundance can be inferred from theoretical solar models by the requirement that the model have the solar luminosity at the age of the Sun. Modern evolution codes give estimates for the initial solar Y in the range $0.26 - 0.28$; the best solar models of Bahcall, Pinsonneault, & Wasserburg (1995) had $Y = 0.272$ and $Y = 0.278$ with and without gravitational settling respectively. A comparison of theoretical stellar models with the Hipparcos main sequence of the Hyades by Perryman et al. (1997) yields $Y = 0.26 \pm 0.02$; for comparison, the solar Y in that study was 0.266 and the solar-scaled helium for the cluster would be 0.28. This agreement between the Sun and Hyades was anticipated and reinforces the notion that stars formed in the current epoch have similar helium abundances.

Nevertheless, we consider what range of Y would be needed to drop the Pleiades main sequence by 0.3 mag, and that value is about $Y = 0.37$. Such a high value of Y for the Pleiades would imply a drastic revision of chemical evolution models and, by extension, would raise the possibility that other clusters might have similar anomalies. MS fitting would therefore require knowledge of both the metal and helium abundances; since helium can only be directly observed in young systems this would make MS fitting unreliable at the 0.3 magnitude level for the majority of clusters.

We believe that this question is best answered by direct measurements of the helium abundance in HII regions and massive stars. We begin with a discussion of the literature on helium abundances; we have also obtained data on the relative helium abundances in the Pleiades and α Per. Neither the field star data nor our Pleiades spectra are consistent with significant variations in the initial helium abundance from the solar value.

Ignoring a deviant few percent of field stars, Nissen’s (1974) study revealed no intrinsic scatter in Y greater than $\sim 10\%$ (compared to the 30-40% deviation required by the Pleiades

stars) in nearby main-sequence field B stars. Gies & Lambert (1992) found helium abundances consistent with both the Sun and the Orion nebula for a sample of 35 B dwarfs; 4 B supergiants in that sample were found to have anomalously high helium abundances. There is evidence that evolutionary effects are responsible for helium enrichment in the most massive stars (see Maeder & Conti 1994, Lyubimkov 1996, Pinsonneault 1997 for reviews), so helium abundances from MS O stars and massive supergiants may not be reliable indicators of the initial Y . The B star field data and the Orion nebula abundances are therefore our best test for the range in helium abundance at solar metal abundance, and they are consistent with only small variations in the initial helium abundance.

For Galactic clusters, however, the picture is less clear. Shipman & Strom (1969), Peterson & Shipman (1973), Nissen (1976), and Lyubimkov (1977) found evidence for 20%-30% variations in Y among young associations, including some systems with significantly lower Y . Lyubimkov suggested an increasing He abundance with *increasing* age amongst the young clusters/associations studied, a conclusion not supported by the subsequent field star work of Gies & Lambert.

Patton (1979) determined He abundances of 60 stars in 8 young clusters and associations. She noted that her initial abundances displayed a range in Y of about 25%, and that this could not be explained by the the usual error sources; she also called attention to a correspondence between He abundance and cluster age. However, Patton shows that binarity may be responsible for observed cluster-to-cluster He abundance dispersions, and the notably low He abundances (observed by others too) seen for a few stars within a given cluster/association. Eliminating *suspected* (but not positively identified) binary systems from her analysis results in cluster He abundances which are identical to within the uncertainties. This highlights the need for secure knowledge of very fundamental stellar parameters (e.g., binarity) before reliable He abundances can be derived.

With this muddled picture of main sequence stellar He abundances, one may wonder if the Pleiades He abundance could be abnormal. Both the Pleiades and α Per are young enough to have B stars, and their helium can be directly measured. The Y values from Lyubimkov (1977) agree to within $\Delta Y \sim 0.015$, which is well within the uncertainties; the Pleiades and α Per Y value is 0.04 larger than the corresponding field star value, but the uncertainties are comparable to this offset. Klochkova & Panchuk (1986) also derived B-star He abundances in both the Pleiades and α Per. They claim to find no difference between the mean abundances that is larger than the uncertainties. However, this conclusion is not clear to us from the abundances listed in their Table II, which do demonstrate quite a very large difference. Unfortunately, only two Pleiades stars are included in the analysis. Therefore small number statistics and the possible effects of binarity make assessment of this difference quite difficult. We attempted a final comparison using the “field” stars from Nissen (1974). This sample includes four α Per stars, and two stars (HR 5191 and 7121) which are suggested members of the purported Pleiades supercluster. The mean Y value is only 0.03 larger for the Pleiades field stars than for the α Per stars; the uncertainties are probably not much smaller than this difference.

To investigate the possibility of a non-standard helium abundance in the Pleiades Fischer & King (1998) observed MS B stars in α Per and Pleiades to differentially compare the helium abundances. Preliminary analysis of the lines strengths for six He lines suggests that the cluster He abundances are identical within an uncertainty of 15%. Any real difference appears to be in the opposite sense of what is needed to make the Pleiades underluminous: the Pleiades line strengths are, if anything, consistently smaller than the α Per counterparts.

4.5. Reddening and Systematic Errors in the Photometry

Reddening will tend to make a cluster MS fainter at a given color. If the reddening is increased the inferred distance modulus will therefore increase. The effect can be roughly estimated as follows : in the color interval that we are using for MS fitting the derivative of M_V with respect to both $B-V$ and $V-I$ is ~ 5 . The extinction $A_V=3.12E(B-V)$ and $E(V-I)_K=1.5E(B-V)$. Adding these effects together an increase in $E(B-V)$ of 0.10 magnitudes would increase V at fixed $B-V$ and fixed $V-I$ by 0.188 (0.5 mag from a shift of 0.1 in $B-V$ - 0.312 mag from extinction) and 0.438 (0.75 mag from a shift of 0.15 in $V-I$ - 0.312 mag from extinction) magnitudes respectively. The relative distances inferred by the two colors can therefore be affected if the reddening is incorrect. In addition the $[\text{Fe}/\text{H}]$ abundances derived for cluster stars are sensitive to T_{eff} , and an increased reddening would imply a higher $[\text{Fe}/\text{H}]$ for a given equivalent width (therefore further increasing the distance modulus). Other colors, such as $R-I$, will be less reddening-sensitive.

Neither the Hyades nor Praesepe show any evidence for reddening along the line of sight; increasing the reddening estimate for the Pleiades would worsen the discrepancy with the Hipparcos distance modulus estimate. Even changing $E(B-V)$ from 0.04 to 0 would only decrease the distance modulus by 0.08 magnitudes. The reddening estimates for the Pleiades have been derived for a wide range of masses and from different techniques; Crawford and Barnes used Stromgren photometry to estimate A_V for B, A, and early F stars in the Pleiades and Praesepe, Prosser and Stauffer used M dwarfs in the same clusters, and Breger used polarization measurements in the Pleiades. We conclude that reddening is not a significant source of uncertainty in distance estimates for the Pleiades. Multicolor distance measurements of the type performed in this paper could be a useful check on the reddening for more heavily obscured systems.

Another possibility is that systematic errors in the photometry could cause errors in

the distance estimates. For the color range that we are considering, the slope of the MS is ~ 5 ; this would require a systematic error of 0.06 magnitudes in $B - V$ to reconcile the Pleiades distance scales, which is unreasonably large. The size of the systematic errors can be constrained by comparing spectroscopic temperature estimates with those based upon colors. In the case of Coma Ber, for example, it appears that spectroscopic temperature estimates are in agreement with the $B - V$ colors of F stars but not with the $V - I$ colors. We note that the slope of the MS in $V - I$ is steeper for F stars than for the cooler stars, and that systematic errors in the $V - I$ photometry might explain the puzzling behavior of Coma Ber. We have attempted whenever possible to rely upon a single source for photometry in a given color for a given cluster. Even in the case of the $V - I$ data we see no evidence of systematic differences between the location on the color magnitude diagram of stars with colors converted to the Cousins system from the Kron system and those converted to the Cousins system from the Johnson system.

For the Pleiades, independent studies (Section 2.2) give consistent photometry for individual stars at the level of the quoted errors (0.01 - 0.02 mag). The 0.3 mag discrepancy between the Hipparcos and MS fitting distance moduli is much too large to be explained by systematic errors in the photometry. High-resolution spectroscopy of the Pleiades is consistent with the observed colors, and the reddening is small. For systems with higher reddening, however, care must be taken when converting between different photometric systems; the Johnson, Cousins, and Kron system I bands have different effective central wavelengths and therefore different reddening corrections.

4.6. Systematic Errors in the Hipparcos Parallaxes

The final possibility is that the Hipparcos Pleiades parallaxes may contain previously undetected systematic errors. If the MS fitting result $m - M = 5.60$ does indeed give the

correct Pleiades distance, then a systematic zero-point error would need to approach the 1 mas level to produce the discordance with the Hipparcos results. Such an error seems impossibly large, in view of the extensive tests (Arenou et al. 1995, 1997) demonstrating the global zero-point error of the Hipparcos parallaxes to be smaller than 0.1 mas. However, global tests have little power to reveal effects occurring on the small angular scale (~ 1 deg) of the Hipparcos spatial correlations (see below). Indeed, the Hipparcos parallaxes of stars in open clusters such as the Pleiades represent the first real opportunity to test for systematic effects on small angular scales. One might well argue that it would only be prudent to consider the Hipparcos cluster results as the first direct tests for small-scale zero-point errors, rather than as cluster distance measurements.

The Hipparcos Pleiades parallax (van Leeuwen & Hansen Ruiz 1997a) is based on measurements of 54 cluster members, ranging in V from 2.8 to 11.5 within 5 deg of the cluster center, so it represents a fairly broad sampling of the cluster. Because Hipparcos observed widely separated (~ 58 deg apart) star fields simultaneously, the parallaxes are inherently on an absolute scale over the whole sky. Over small regions of the sky ($\lesssim 2$ deg), however, the astrometric results are positively correlated because neighboring stars (within the $0.9 \text{ deg} \times 0.9 \text{ deg}$ Hipparcos field of view) tended to be observed on the same great circles the satellite swept out over the sky (Lindgren 1988, 1989). A comprehensive discussion of the Hipparcos mission and data reductions is given in Volumes 1–3 of the Hipparcos Catalogue (ESA 1997). The spatial correlations may significantly impact the astrometric results for star clusters, whose angular size is of the same order as the Hipparcos correlation scale. To account for this, van Leeuwen & Hansen Ruiz (1997a) re-calculated the Pleiades mean parallax from the intermediate Hipparcos data. For this paper, one of us (R.B.H.) has re-examined the individual Pleiades parallaxes from the Hipparcos Catalogue.

Moreover, besides the spatial correlations, there is a different type of correlation

affecting the Hipparcos results – the statistical correlations among the five astrometric parameters (position, proper motion, and parallax), arising from the imperfect distribution of Hipparcos observations on the sky over time.

In classical parallax work (cf. Vasilevskis 1975), the time distribution of observations over a star’s parallactic ellipse is controlled to maximize the parallax factors and minimize the correlations between position, proper motion, and parallax. This is easy to achieve from the ground, but Hipparcos could not do this because of the limited span of observations and the pattern of scans of the sky, as explained in Section 3.2.4 (pp. 321-325) of the Hipparcos Introduction (ESA 1997, Vol. 1). Figures 3.2.42 to 3.2.61 of that work illustrate the patterns of the correlations over the sky; Figure 3.2.66 (p. 363) shows histograms of the 10 correlations. The RMS values are ~ 0.2 , and large areas of the sky show correlations averaging 0.4 or more in size. It must be emphasized that these correlations are substantially larger than would be considered acceptable in ground-based parallax observations.

For parallax work, the most important correlation is ρ_{α}^{π} , between parallax and right ascension (Field H20 in the Hipparcos Catalogue). This is because, over most of the sky, most of the extent of the parallactic ellipse is in right ascension. The Hipparcos ρ_{α}^{π} correlation is shown in Fig. 3.2.44 of the Hipparcos Introduction. Large values of ρ_{α}^{π} were caused in certain areas of the sky by the unfortunate circumstance of unequal observations on both sides of the Sun, as discussed on p. 325 of the Hipparcos Introduction.

This happens to impact the Pleiades particularly badly. The mean value of ρ_{α}^{π} near the Pleiades center is +0.4; this is at the 96th percentile in the histogram in Fig 3.2.66. The question this raises is whether this large correlation, caused by the time distribution of Hipparcos observations of the Pleiades stars, has any effect on the parallax values.

We stress again that this is a different effect from the spatial correlation that exists because Hipparcos astrometric data over small (~ 1 deg) areas of the sky are not fully

independent measurements.

In Figure 19 we plot parallax vs. the correlation ρ_α^π for 49 Pleiades members verified by proper motion, radial velocity, and position in the color-magnitude diagram. (Mermilliod et al’s 51 stars and van Leeuwen et al’s 54 are virtually the same set as these; we rejected several additional stars on account of problems noted in Fields H30 and H59 of the Hipparcos Catalogue.) This plot shows several interesting things.

The filled symbols are 12 bright ($V < 7$) stars within ~ 1 deg of the cluster center with correlations $\rho_\alpha^\pi \geq +0.34$ (the mean value for the whole sample). Due to the spatial correlation effect, these 12 stars all have nearly the same parallax (mean 8.86 mas, RMS dispersion 0.45 mas; χ^2 too small at the 0.995 significance level). Because Hipparcos’ errors are smallest for bright stars, these stars carry much of the weight of the Pleiades parallax.

There is a clear trend (slope) of parallax vs. ρ_α^π correlation; a weighted least-squares solution gives a slope of $+3.04 \pm 1.36$ mas per unit correlation. The solid line in Fig. 19 is this slope, run through the mean point (+0.34,+8.53). The dashed lines show $\pm 1\sigma$ slopes. The intercept at zero correlation is $\pi = 7.49 \pm 0.50$ mas, quite consistent with the MS fitting distance.

Figure 20 plots parallax vs. distance from the cluster center. The filled symbols are the same 12 bright stars with high ρ_α^π as in Fig. 16. The open symbols are the 15 stars with $\rho_\alpha^\pi < +0.25$, with no restriction on magnitude or distance. The two sets of stars barely overlap because the brightest stars in the Pleiades are highly concentrated to the cluster center. The low-correlation stars lie farther from the Pleiades center and show a much larger parallax scatter, reflecting (a) the larger errors for fainter stars and (b) the lack of spatial correlations on scales $\gtrsim 1$ deg. Moreover, their mean parallax is smaller (reflecting the slope discussed above). For the 15 stars with $\rho_\alpha^\pi < +0.25$, the weighted mean parallax is 7.46 ± 0.43 mas. The RMS dispersion is 1.66 mas, consistent with the published parallax

errors.

This exercise is not intended to be a definitive re-determination of the Pleiades parallax; that would require going back to the intermediate Hipparcos data as per van Leeuwen et al (1997), and exploring the effects of both the ρ_α^π and the spatial correlations at that level. However, it is quite clear that (a) small-angular-scale systematic effects at the 1 mas level are present in the Hipparcos Pleiades parallaxes; (b) these effects are related to the high values of the ρ_α^π correlation near the cluster center; (c) the bright stars within ~ 1 deg of the center, which carry most of the weight of the mean parallax, are the most severely affected; and (d) the stars with lower ρ_α^π correlations, far enough ($\gtrsim 1$ deg) from the center to be unaffected by the spatial correlation, have smaller parallaxes, consistent with the MS fitting distance.

We also looked for effects of the ρ_α^π correlation in the Hyades, Praesepe, α Per, and Coma Ber clusters. In Figures 21–24 we present the parallax vs. correlation plots for those clusters. The Hyades, Praesepe, and α Per clusters also have large values of ρ_α^π , but the slope ($d\pi/d\rho$) present in the Pleiades data does not occur in these clusters, where the MS fitting distances and the Hipparcos distances are in good agreement. The data for Coma Ber do show a slope $d\pi/d\rho = -4.0 \pm 2.1$ mas, but the range of ρ_α^π is small, and the mean is near zero.

5. Resolution of the Problem

The Hipparcos distances to the open clusters can be regarded as either a test of the theory of stellar structure and evolution or as a test of the parallaxes themselves. To distinguish between the two it is necessary to determine what the errors in stellar interiors-based cluster distances are. We have performed a detailed multicolor analysis of

the distances to the nearby open clusters, and verify that MS fitting can be performed to a precision of order 0.05 magnitudes. With the exception of Coma Ber, distance estimates from $B-V$ and $V-I$ colors can be used to get photometric $[\text{Fe}/\text{H}]$ values accurate at the 0.05 dex level, and these estimates are in good agreement with those obtained from high-resolution spectroscopy. There is a small zero-point shift, of order 0.04 dex, between our photometric abundance scale and that of Friel and Boesgaard; if we adopted our zero-point the distance estimates we have reported would all be increased by 0.04 magnitudes. We also note that the distances inferred for rapid rotators are not consistent for the two colors; this implies that color temperatures for these stars may be in error, especially if they are derived from $B-V$ colors. This may play some role in the lithium-rotation correlation seen in young rapidly rotating stars.

We have shown that the internal consistency of MS fitting is high and, in the particular case of the open clusters, the systematic errors are small. The basic cluster data (abundances, reddening, etc.) are also well established for the systems that we have studied in this paper. The extremely good agreement between helioseismology and theoretical solar models places strong constraints on missing physics in the models, and by extension the properties of solar analogs should be accurately represented by the models. For all of these reasons we believe that the open cluster distance scale from MS fitting is on very strong ground.

The Hipparcos mission permits a comparison of parallax and MS fitting distances for a number of open clusters. In two of the systems that we have studied (α Per and the Hyades) the two distance scales are in very good agreement. In Coma and the Pleiades they disagree at the 0.2 and 0.3 magnitude level respectively; these differences are at the 3.4 and 3.7 σ level respectively. The different distance scales for Praesepe are either very close (0.08 mag) or discrepant (0.33 mag) depending upon which of the Hipparcos distance

measurements is adopted; the latter would be a 2σ disagreement. We have searched for sources of error in the MS fitting distances of Coma Ber and the Pleiades. The $V - I$ photometry of Coma yields a distance that disagrees both with Hipparcos and $B - V$; this can be traced to a discrepancy in the temperature scales for the two colors in this cluster. Although we believe that there are a number of indications that the $B - V$ temperature scale is correct (consistency with spectroscopic temperatures and Stromgren photometry, for example) reobserving this cluster in IR and near-IR colors would be highly desirable to quantify the magnitude of the problem.

In the case of the Pleiades there is no such ambiguity; different colors yield identical distances. Errors in the metal abundance and reddening as a solution can be rejected on a variety of grounds. The increase in the cluster helium abundance needed to reconcile the distance estimates is large and not consistent with direct measurements. Furthermore, we can find no counterparts of the Pleiades in the field, i.e. intrinsically faint solar abundance stars (Soderblom et al. 1998). We are therefore left with the uncomfortable choices of either requiring unknown physics in the interiors models or a problem with the Hipparcos parallax distance scale to the Pleiades. The former choice is made even less attractive by the requirement that the models retain agreement with the Sun, the other clusters, and numerous other tests of the theory of stellar structure and evolution. We therefore believe that the latter explanation is more likely.

We have shown that there is evidence in the Pleiades data for systematic errors in the parallaxes on small angular scales. The same trends are not present in the clusters where the two distance scales agree; they may also be present in the Coma Ber cluster. Clusters such as the Pleiades provide many more stars within a small region of the sky than are present for the sky as a whole, and they are therefore uniquely suited to test systematic effects at small angular scales. The other clusters and the Pleiades show no evidence for

systematic errors on scales larger than 1 deg. The Pleiades results suggest that individual parallax measurements with large ρ_{α}^{π} correlations should be treated with caution. The implications of this result for other applications of the Hipparcos parallaxes will depend upon the characteristics of the sample. For large samples over large regions of the sky the net effect will be a modest increase in the overall error. A numerical example: Arenou et al. (1997, pp.441-443) find the overall ratio of Hipparcos “external” to “internal” errors to be 1.06 ± 0.07 from clusters and 1.04 ± 0.04 from distant stars. With the internal error ~ 1 mas, this is equivalent to an additional (in quadrature) error ~ 0.2 - 0.4 mas. This may in fact be the RMS size of the small-scale errors.

D.R.S. acknowledges partial support from NASA grant NAGW-4837. R.B.H acknowledges partial support from NASA Grant NAG5-4830 and NSF grant AST 9530632. J.R.S. acknowledges partial support from NASA grants NAGW-2698 and NAG5-3363. We wish to thank Debra Fischer, Eileen Friel, Burt Jones, Jean-Claude Mermilliod, and Donald Terndrup for helpful discussions.

REFERENCES

- Allard, F., Hauschildt, P.H., Alexander, D.R., and Stafffield, S. 1997, *ARA&A*, 35, 137
- Allen, C.W. 1973 *Astrophysical Quantities*, Third Edition (London, Athlone Press)
- Alonso, A., Arribas, S., & Martinez-Roger, C. 1996, *A&A*313, 873
- Arenou, F., Lindegren, L., Froeschle, M., Gomez, A.E., Turon, C., Perryman, M.A.C. & Wielen, R. 1995, *A&A*, 304, 52
- Arenou, F., Mignard, F. & Palasi, J. 1997, Chapter 20, Verification of Parallaxes, in the *Hipparcos and Tycho Catalogues*, Vol. 3, Construction of the Hipparcos Catalogue, ESA SP-1200
- Bahcall, J.N., Pinsonneault, M.H., & Wasserburg, G. 1995, *Rev. Mod. Phys.*, 67, 781
- Balachandran, S., Lambert, D. L., & Stauffer, J. R. 1988, *ApJ*, 333, 267
- Bessell, M. S. 1979, *PASP*, 91, 589
- Bessell, M. S. & Brett, J.M. 1988, *PASP*, 100, 1134
- Bessell, M. S. & Weis, E.W. 1987, *PASP*, 99, 642
- Boesgaard, A. M., Budge, K. G., & Ramsay, M. E. 1988, *ApJ*, 327, 389
- Boesgaard, A. M., 1987, *ApJ*, 321, 967
- Boesgaard, A. M., 1989, *ApJ*, 336, 798
- Boesgaard, A. M., & Friel, E. D. 1990, *ApJ*, 351, 467
- Bouvier, J., Rigaut, F., & Nadeau, D. 1997, *A&A*, 323, 139
- Breger, M. 1986, *ApJ*, 309, 311
- Cayrel, R., Cayrel de Strobel, G., & Campbell, B. 1988, *IAU Symposium* 132, 449
- Chaboyer, B., Demarque, P., Kernan, P.J., & Krauss, L.M. 1998, *ApJ*, 494, 96

- Crawford, D. & Barnes, J. 1969, AJ, 74, 818
- Crawford, D. & Barnes, J. 1974, AJ, 79, 687
- Crawford, D. & Perry, C. 1966, AJ, 71, 206
- Crawford, D. & Perry, C. 1976, AJ, 81, 419
- Eggen, O. J. 1986, PASP, 98, 755
- ESA 1997, The Hipparcos and Tycho Catalogues, ESA SP-1200
- Fischer, D. & King, J. 1998, in preparation
- Friel, E. D., & Boesgaard, A. M. 1992, ApJ, 387, 170
- Giannuzzi, M. A. 1995, A&A, 293, 360
- Gies, D.R. & Lambert, D.L. 1992, ApJ, 387, 673
- Gratton, R.G., Fusi Pecci, F., Carretta, E., Clementini, G., Corsi, C.O., & Lattanzi, M.
1997, ApJ, 491, 749
- Green, E.M. 1988, in Calibration of Stellar Ages, ed. A.G.D. Phillip (Schenectady,
NY:Davis), 81
- Griffin, R.F., Gunn, J.E., Zimmerman, B.A., & Griffin, R.E.M. 1988, AJ, 96, 172
- Johnson, H.J. 1952, ApJ, 116, 640
- Johnson, H.J. & Knuckles, C.F. 1955, ApJ, 122, 209
- Johnson, H.J. & Mitchell, R.I. 1958, ApJ, 128, 31
- King, J.R. 1993, PhD thesis, University of Texas at Austin.
- King, J. R., Soderblom, D. R., Fischer, D., & Jones, B.F. 1997, in preparation
- King, J. R., & Hiltgen 1996, PASP, 108, 246
- Klochova, V. G., & Panchuk, V. E. 1986, Sov. Astr. Let., 12, 387

- Kurucz, R.L. 1991a, private communication
- Kurucz, R.L. 1991b, private communication
- Landolt, A. 1979, ApJ, 231, 468
- Lindgren, L. 1988, in Torra, J. and Turon, C., Scientific Aspects of the Input Catalogue Preparation II, Proc. Coll. Sitges (Catalonia), 179
- Lindgren, L. 1989, in Perryman et al. 1989, The Hipparcos Mission, ESA SP-1111, Vol. III, 311
- Lyubimkov, L.S. 1977, Astrofizika, 13, 139
- Lyubimkov, L.S. 1996, Ap&SS, 243, 329
- Maeder, A. & Conti, P.S. 1994, ARA&A, 32, 227
- McWilliams, A. 1997, ARA&A, 35, 503
- Mendoza, E.E. 1967, 4, 149
- Mermilliod, J.-C., Rosvick, J.M., Duquennoy, A.M., & Mayor, M. 1992, A&A, 265, 513
- Mermilliod, J.-C., Turon, C., Robichon, N., Arenou, F., & Lebreton, Y. 1997, in *Hipparcos Venice '97*, eds. B. Battick & M. A. C. Perryman (ESA), 643
- Mitchell, R.I. 1960, ApJ, 132, 68
- Nissen, P. E. 1974, A&A, 36, 57
- Nissen, P. E. 1976, A&A, 50, 343
- Patton, G. Y. 1979, Ph. D. Dissertation, SUNY-Stony Brook
- Perryman, M. A. C., Brown, A.G.A., Lebreton, Y., Gomez, A., Turon, C., Cayrel de Strobel, G., Mermilliod, J.-C., Robichon, N., Kovalevsky, J., & Crifo, F. 1997, A&A, in press
- Peterson, D. M., & Shipman, H. L. 1973, ApJ, 180, 635

- Pinsonneault, M.H. 1997, ARA&A, 35, 557
- Pont, M., Mayor, M., Turon, C., & Vandenberg, D.A. 1998, A&A, 329, 87
- Prosser, C.F. 1992, AJ, 103, 488
- Prosser, C.F. 1994a, AJ, 103, 488
- Prosser, C.F. 1994b, AJ, 107, 1422
- Reid, I.N. 1997, AJ, 114, 161
- Robichon, N., Arenou, F., Turon, C., Mermilliod, J.-C., & Lebreton, Y. 1997, in *Hipparcos Venice '97*, eds. B. Battick & M. A. C. Perryman (ESA), 567
- Rogers, F.J., & Iglesias, C.A. 1992, ApJ, 397, 717
- Rogers, F.J., Swenson, F.J., & Iglesias, C.A. 1996, ApJ, 456, 902
- Saumon, D., Chabrier, G., & van Horn, H.M. 1995, ApJS, 99, 713
- Schwan, H. 1991, A&A, 243, 386
- Shipman, H. L., & Strom, S. E. 1969, BAAS, 1, 362
- Soderblom, D.R., Stauffer, J. R., Hudon, J. D., & Jones, B. F. 1993, ApJS, 85, 315
- Soderblom, D.R., King, J.R., Hanson, R.B., Jones, B.F., Fischer, D., Stauffer, J.R., & Pinsonneault, M.H. 1998, ApJ, in press
- Stauffer, J.R. 1980, AJ, 85, 1341.
- Stauffer, J.R. 1982a, AJ, 87 1507
- Stauffer, J.R. 1982b, PASP, 94, 678
- Stauffer, J.R. 1984, ApJ, 280 189
- Stauffer, J.R. & Hartmann, L. 1987, ApJ, 318, 337
- Stauffer et al. 1985, ApJ, 289, 247

- Stauffer et al. 1989, ApJ, 346, 160
- Taylor, B.J. 1994, PASP, 106, 600
- Upgren, A.R., Weis, E.W., and DeLuca, E.E. 1979, AJ, 84, 1586.
- VandenBerg, D. A., & Bridges, T. J. 1984, ApJ, 278, 679
- van Leeuwen, F., & Hansen Ruiz, C. S. 1997a, in *Hipparcos Venice '97*, eds. B. Battick & M. A. C. Perryman (ESA), 689
- van Leeuwen, F., & Hansen Ruiz, C. S. 1997b, A&AS, in press
- Vasilevskis, S. 1975, Publ. Lick Obs. 22, Part IV
- Weis, E.W. 1981, PASP, 93, 437
- Wyse, R.F.G. & Gilmore, G. 1995, AJ, 110, 2771

Fig. 1.— The Pleiades compared with Praesepe (top panel) and α Per (bottom panel) in $B-V$. Stars have been shifted by the Hipparcos distances in Table 1; photometry sources are listed in Section 2. The Pleiades stars are the filled symbols and stars in the other clusters are the open symbols.

Fig. 2.— The Pleiades compared with Praesepe (top panel) and α Per (bottom panel) in $(V-I)_C$. Stars have been shifted by the Hipparcos distances in Table 1; Johnson and Kron $V-I$ have been converted to Cousins $V-I$ as described in Section 2. Photometry sources are listed in Section 2. The Pleiades stars are the filled symbols and stars in the other clusters are the open symbols.

Fig. 3.— M_V for a 1 Gyr as a function of $\log T_{eff}$ (top panel), $B-V$ (middle panel) and $(V-I)_C$ (bottom panel). In each panel the bottom, middle, and top solid lines are $[\text{Fe}/\text{H}]=-0.3, 0$, and $+0.2$ respectively; the dashed line is a $[\text{Fe}/\text{H}] = 0$ isochrone with a helium abundance of 0.37 (0.1 higher than the solar calibrated value).

Fig. 4.— Isochrones with an age of 1 Gyr and a range of $[\text{Fe}/\text{H}]$ are compared with field stars with $M_V > 4$ and relative parallax errors less than 5% in $B-V$ (top panel) and $(V-I)_C$ (bottom panel). Age effects are responsible for the departure of the data from the isochrone for $M_V < 5$. The top, middle, and bottom isochrones are respectively $[\text{Fe}/\text{H}]=+0.2, 0$, and -0.3 .

Fig. 5.— Theoretical 600 Myr isochrones in the $M_{bol}/\log T_{eff}$ plane for solar $[\text{Fe}/\text{H}]$ (bottom lines) and Hyades $[\text{Fe}/\text{H}]$ (top lines). The solid lines are from this paper and the dashed lines are from Perryman et al. 1997.

Fig. 6.— M_V as a function of $(B-V)$ (top panel) and $(V-I)_C$ (bottom panel) for Hyades members with Hipparcos parallax measurements and $(B-V) > 0.5$. Photometry sources are listed in Section 2. Solid points are single stars; open points are binaries from Griffin et al.

1988. The empirical fits to the single star MS used in this paper are the solid lines.

Fig. 7.— The difference between the Hipparcos M_V and that predicted from the isochrones for single Hyades stars with parallax measurements is plotted as a function of $(B-V)$ (top panel) and $(V-I)_C$ (bottom panel). The solid lines indicates the corrections as a function of color that were applied to the isochrones for consistency with the shape of the Hyades MS.

Fig. 8.— Histogram of differences between the Hipparcos M_V and that predicted from the isochrones for $(B-V)$ (top) and $(V-I)_C$ (bottom). Data are binned in 0.1 magnitude intervals; single stars are the dark bins and binaries are the light bins.

Fig. 9.— Histogram of distance modulus estimates for Pleiades members, binned in 0.05 magnitude intervals. The isochrones were used for the top panels, while the isochrones were corrected to the shape of the Hyades MS in the bottom panels; those on the left side are distances from the $(B-V)$ color and those on the right side are distances from the $(V-I)$ color. that predicted from the isochrones for $(B-V)$ (top) and $(V-I)_C$ (bottom). Stars with $B-V$ from 0.5 – 0.75 are the dark bins and stars with $B-V$ from 0.76 – 0.9 are the light bins.

Fig. 10.— 100 Myr theoretical isochrones with $[\text{Fe}/\text{H}]=-0.03$ shifted to a distance of 5.60 are compared with the Pleiades in $(B-V)$ (top) and $(V-I)$ (bottom). Single stars are the filled squares, open circles are binaries from Bouvier, Rigaut, & Nadeau 1997, and open circles with a cross are rapid rotators. The solid lines are the isochrones and the dashed lines are the same isochrones with the Hyades MS shape.

Fig. 11.— As the top two panels of Figure 9, but with a wider color range ($B-V$ from 0.5 – 1.0) and with data binned in 0.1 magnitude intervals. Single stars are the dark bins, binaries from Bouvier, Rigaut, & Nadeau 1997 are the light bins, and rapid rotators are the

striped bins.

Fig. 12.— As for Figure 9, except for Praesepe relative to a 600 Myr isochrone.

Fig. 13.— 600 Myr theoretical isochrones with $[\text{Fe}/\text{H}]=+0.04$ shifted to a distance of 6.16 are compared with Praesepe in $(B-V)$ (top) and $(V-I)$ (bottom). The solid lines are the isochrones and the dashed lines are the same isochrones with the Hyades MS shape.

Fig. 14.— Histogram of distance modulus estimates for α Per members, binned in 0.05 magnitude intervals. The top panel shows distances from the $(B-V)$ color and those on the bottom panel are distances from the $(V-I)$ color; isochrones with the Hyades MS shape yield similar results and are not shown. Slow rotators are the dark bins and rapid rotators ($v \sin i > 49$ km/s) are the light bins.

Fig. 15.— 50 Myr theoretical isochrones with $[\text{Fe}/\text{H}]=-0.05$ (solid lines) shifted to a distance of 6.23 are compared with α Per in $(B-V)$ (top) and $(V-I)$ (bottom). Slow rotators are the filled squares and rapid rotators are the open circles with a cross.

Fig. 16.— Histogram of distance modulus estimates for Coma Ber members, binned in 0.05 magnitude intervals. The top panel shows distances from the $(B-V)$ color and those on the bottom panel are distances from the $(V-I)$ color. Stars with $(B-V)$ from 0.5 – 0.9 are the dark bins, which corresponds to the color interval used for the other clusters. The light bins are hotter stars with $(B-V)$ from 0.35 – 0.499. Note that for the cooler stars there is no well-defined peak for $(V-I)$, and that the overall peak disagrees significantly with the Hipparcos and $(B-V)$ distances (see text).

Fig. 17.— Effective temperatures for Coma Ber members from Boesgaard (1987) compared with those predicted from $(B-V)$ (top) and $(V-I)$ (bottom).

Fig. 18.— 500 Myr theoretical isochrones (solid lines) with $[\text{Fe}/\text{H}]=-0.07$ shifted to a distance

of 4.54 are compared with Coma Ber in $(B-V)$ (top) and $(V-I)$ (bottom).

Fig. 19.— Hipparcos parallax vs. the correlation ρ_α^π for 49 Pleiades members. Filled symbols are 12 bright ($V < 7$) stars within ~ 1 deg of the cluster center with correlations $\rho_\alpha^\pi \geq +0.34$. Vertical dotted lines mark $\rho_\alpha^\pi = 0$ and the mean value $+0.34$. Sloping lines represent the weighted least-squares relation $\pi = 8.53 + (3.04 \pm 1.36)(\rho_\alpha^\pi - 0.34)$ mas.

Fig. 20.— Hipparcos parallax vs. angular distance from the Pleiades cluster center. The filled symbols are the same 12 bright stars with high ρ_α^π as in Fig. 14. The open symbols are the 15 stars with $\rho_\alpha^\pi < +0.25$, with no restriction on magnitude or distance. The long-dashed line marks the mean parallax (7.46 mas) for these 15 stars. The dotted line marks the mean parallax (8.86 mas) for the 12 bright stars.

Fig. 21.— Hipparcos parallax vs. the correlation ρ_α^π for the 40 Hyades cluster members in Table 8 of Perryman et al. (1997). Vertical dotted lines mark $\rho_\alpha^\pi = 0$ and the mean value $+0.29$. Horizontal dotted line marks the mean parallax 22.05 mas.

Fig. 22.— Hipparcos parallax vs. the correlation ρ_α^π for 20 Praesepe cluster members verified by proper motion and position in the color-magnitude diagram. Vertical dotted lines mark $\rho_\alpha^\pi = 0$ and the mean value $+0.27$. Horizontal dotted line marks the mean parallax 5.72 mas.

Fig. 23.— Hipparcos parallax vs. the correlation ρ_α^π for 51 α Per cluster members verified by proper motion, radial velocity, and position in the color-magnitude diagram. Vertical dotted lines mark $\rho_\alpha^\pi = 0$ and the mean value $+0.23$. Horizontal dotted line marks the mean parallax 5.48 mas.

Fig. 24.— Hipparcos parallax vs. the correlation ρ_α^π for 26 Coma Ber cluster members verified by proper motion and position in the color-magnitude diagram. Vertical dotted lines mark $\rho_\alpha^\pi = 0$ and the mean value -0.04 . Horizontal dotted line marks the mean parallax 11.41

mas.

Figure 1




Figure 2





Figure 3




Figure 4





Figure 5




Figure 6





Figure 7







Figure 8









Figure 10







Figure 11









Figure 13





Figure 14





Figure 15





Figure 16





Figure 17





Figure 18

